

特別講演

ビキニ被災者20年間の臨床的、特に血液学的観察

熊 取 敏 之

放射線医学総合研究所障害臨床研究部

ACTA HAEM. JAP. 38: 635-645, 1975

Clinical, especially Hematological Observations over the 20-Year Period
on the Japanese Fisherman Exposed to Fallout in 1954

Toshiyuki KUMATORI

Division of Radiation Health, National Institute of Radiological Sciences, Chiba

On March 1 in 1954, 23 Japanese fishermen aged from 18 to 39 were exposed to radioactive fallout produced by the thermonuclear test explosion at Bikini Lagoon. After 14 days navigation they returned to their harbor where they were found to have been injured by radioactive materials. They were hospitalized for 13 months. After being discharged most of them have been examined so far as possible on an annual basis.

The fishermen were irradiated in the following three ways: (1) From the radioactive materials adhered to the skin. (2) Externally from the radioactive materials in the cabins, on the deck etc.. (3) Internally from the radioactive materials entered various organs. While the estimate of radiation dose to skin as well as the dose by internal radiation were difficult, the estimated external radiation dose was approximately 170-500 rad for 14 days, about, half or more of which was irradiated on the first day. The dose to each person differed depending on his behavior on the boat and the position of his cabin, and had the intimate relationship with his minimum value of leukocytes and neutrophils. Soon after exposure most of the fishermen experienced anorexia, fatigue, and conjunctivitis, and in some of them nausea and vomiting occurred.

Shortly after the exposure, they suffered from erythema which was followed by edema, vesicle, erosion, ulceration or necrosis. Epilations were observed in 20 cases. Though the skin injuries recovered gradually, some persons continued to show residual changes. However, neither chronic radiation dermatitis nor malignant changes was observed.

Leukocytes count decreased gradually, showing minimum count at 4th-7th week. 5 cases revealed a count of less than 2,000/mm³, 13 less than 3,000 and 5 less than 4,000. At first lymphopenia was noticed and then neutropenia became marked. At the recovering stage, eosinophilia and the appearance of immature neutrophils were observed in many cases. In severe cases slight anemia was observed, accompanied by the depression of reticulocytes. Platelets count showed minimum value at the 4th-7th week. Slight hemorrhagic tendencies were observed in severe cases. Bone marrow of severe cases was highly hypoplastic at the critical stage, which changed to slightly hypoplastic and turned into almost normal marrow. The coexistence of hypoplastic area and hyperplastic area was observed in histological sections at the recovery stage and even in the examinations of 10-15 years after the exposure. Morphological abnormalities of blood cells were noticed in both peripheral blood and bone marrow. The cumulative distribution curve of leukocytes counts displaced slightly to the left of normal one even 6 years after the exposure.

Follow-up of chromosome observations in blood cells has been performed since 1964. Even 20 years after exposure, cells with chromosome abnormalities (both Cu and Cs cells) exist in the peripheral lymphocytes with much higher frequencies than in general population. While the frequency of Cu cells (dicentric and rings) was decreasing Cs cells remained fairly constant. The frequencies of

Reprinted from "Biological Aspects
of Radiation Protection"
edited by T. Sugahara and O. Hug,
IGAKU SHOIN LTD., Tokyo, 1971.

HEMATOLOGICAL EFFECTS ON HEAVILY IRRADIATED JAPANESE FISHERMEN

TOSHIYUKI KUMATORI

*Division of Radiation Health,
National Institute
of Radiological Sciences
Chiba, Japan*

1971



IGAKU SHOIN LTD. TOKYO

HEMATOLOGICAL EFFECTS ON HEAVILY IRRADIATED JAPANESE FISHERMEN

TOSHIYUKI KUMATORI

*Division of Radiation Health,
National Institute
of Radiological Sciences
Chiba, Japan*

INTRODUCTION

On 1 March in 1954, a thermonuclear test explosion was performed by the United States Authorities at Bikini Lagoon in the Pacific Ocean. At that time one Japanese fishing boat "the 5th Fukuryu-maru" was located at about 90 miles east from explosion center. The boat was nearly 100 tons and 23 fishermen aged from 18 to 39 were on board.

At about 3:40 a. m., while they were occupied in fishing tuna they saw a huge red light in the west and heard a detonation several minutes later. At about 7:00 a. m. white material began to fall, which continued for approximately 5 hours. They gave up fishing and returned to their mother port on 14 March 1954.

After landing, all the fishermen were found to have been injured by the radioactive materials. Seven of them were hospitalized to the Tokyo University Hospital and the other 16 were received by the First National Hospital of Tokyo by 28 March. They were discharged from both hospitals in May 1955, except for one fatal case who died on 23 September 1954. After being discharged we continued the follow-up studies so far as possible on annual basis. However, mainly because of the varied status of their occupations and widely distributed addresses, it was not always possible to get them all together for medical examination.

Medical data of the fishermen have been reported several times (MIYOSHI and KUMATORI, 1955; KOYAMA et al., 1955; MIKAMO et al., 1956; MIYOSHI and KUMATORI, 1962; KUMATORI and MIYOSHI, 1963; MIYOSHI and KUMATORI, 1964; KUMATORI et al., 1965).

FALLING OF RADIOACTIVE MATERIALS

During the falling of the white materials, when its intensity was greatest, these persons were unable to keep their mouths and eyes open. Their footprints were clearly marked on the deck covered by fallout.

The fishermen were in
(1) From the radioact
(2) Externally from th
the deck etc.
(3) Internally from th

The diameters of the reef fragments, ranged from 1954). According to the March, rare-earth element activity and uranium concentration specific activity of the material of these data, a value of 1 a. m. on 1 March. The internally deposited radio

However, radiation lesions. External radiation exposure dose for 14 days cent of these doses was based on the results of the exponential decay curves of many

Table 1. Exposure

Subject No.

T-1
T-2
T-3
T-5
T-6
T-7
T-8
K-1
K-2
K-3
K-4
K-5
K-6
K-7
K-8
K-9
K-10
K-11
K-12
K-13
K-14
K-15
K-16

* T-3 put the fall-out added in total.

HEAVILY FISHERMEN

STATE OF IRRADIATION AND ESTIMATED RADIATION DOSE

The fishermen were irradiated in the following 3 ways:

- (1) From the radioactive materials adhering to the body surface.
- (2) Externally from the radioactive materials deposited in the cabins, on the deck etc.
- (3) Internally from the radioactive materials entering various organs.

The diameters of the fallout materials, which were the powders of coral-reef fragments, ranged from 19 to 460 μ , mainly 100 to 400 μ (KIKUCHI et al., 1954). According to the results of radiochemical analysis of fallout on 26 March, rare-earth elements contributed about 50 per cent of the total radioactivity and uranium contributed about 20 per cent (KIMURA et al., 1956). The specific activity of the material was 0.37 mCi/g on 23 April. By extrapolation of these data, a value of 1.4 Ci/g was obtained as the specific activity at 7:00 a.m. on 1 March. The estimation of radiation dose, especially that from internally deposited radioactive materials was difficult.

However, radiation from body surface was large enough to cause skin lesions. External radiation dose was estimated as shown in Table 1. The exposure dose for 14 days ranged from about 170 to 690 R, and nearly 60 per cent of these doses was received on the first day. This estimation was based on the results of the experimental reproduction of ash-fall, extrapolation of decay curves of many places of the boat, and on detailed investigation of

Table 1 Estimated dose of whole body gamma radiation.

Subject No.	First Day	Total
T-1	240~290 R	450~500 R
T-2	210~260	390~440
T-3	150~200	260~310* (360~410)
T-5	400~430	660~690
T-6	130~180	200~250
T-7	140~190	220~270
T-8	310~360	520~570
K-1	190~220	310~340
K-2	130~180	200~250
K-3	140~190	230~280
K-4	120~170	190~240
K-5	140~190	220~270
K-6	180~230	300~350
K-7	230~280	340~390
K-8	220~270	380~430
K-9	310~360	550~600
K-10	140~190	230~280
K-11	120~170	170~220
K-12	100~150	170~220
K-13	250~300	370~420
K-14	420~500	510~590
K-15	140~190	210~260
K-16	120~170	190~240

* T-3 put the fall-out material close to his bed. Therefore about 100 R should be added in total.

ion was performed by Pacific Ocean. At that u" was located at about nearly 100 tons and 23

a fishing tuna they saw n several minutes later. ich continued for ap- turned to their mother

ave been injured by the l to the Tokyo Univer- First National Hospital both hospitals in May nber 1954. After being as possible on annual of their occupations and ble to get them all to-

l several times (MIYOSHI al., 1956; MIYOSHI and r and KUMATORI, 1964;

TRIALS

s intensity was greatest, ys open. Their foot- illout.

each patient's behavior on the boat. The dose to each person differed depending on his behavior and the position of his cabin.

The integrated dose to the thyroid from ^{131}I was inferred as about 20 to 120 rads from external countings of radioactivity in the thyroid region during the 4 to 7 weeks after the initial exposure.

Significant amounts of radioactivity were found in the urine samples collected at about 4 weeks after the explosion. The radioactivity decreased rapidly, namely at about 6 months post explosion the activity in the urine was barely detectable.

Radiochemical analysis of several organs from the fatal case showed that the radioactivity of these organs was clearly higher, when compared with the controls, although it was not extremely high (KIMURA et al., 1956).

After $8\frac{1}{2}$ years and 10 years, ^{137}Cs - and ^{90}Sr content in the urine of the fishermen revealed no significant increase. At the same time the results of whole body counting showed no significant difference between fishermen and controls.

Thus, the external irradiation seemed to play an important role in early effects.

SYMPTOMS AND SIGNS IN THE EARLY STAGE

Figure 1 summarizes the symptoms and signs in the early stage.




Fig. 1 Symptoms and signs in the early stage.

HEMA

1. Leukocytes

The total number of counts at about the 4th to count of less than 2,000/ μl one case, the leukocyte between these minimum Similar relationship was the dose as shown in Fi

Fig. 2 Correlation between the estimated doses to the whole body and the minimal number of neutrophils.

In most cases a cells increased in ea some cases, metamy especially with the damage, a considerable end stage. Lympho cases. Leukocyte eosinophilia was re Eosinophilia continued and one case still r

2. Erythrocytes

At the time of was observed in th The Price-Jones c of normal at first, a corresponded to c

3. Platelets, etc.

Platelet count Most of the fish of other tests rel in general and r

1. Comparison v

When these l

HEMATOLOGICAL OBSERVATIONS

1. Leukocytes

The total number of leukocytes decreased gradually, showing minimum counts at about the 4th to 8th week after the exposure. Five cases revealed a count of less than 2,000/mm³, 13 less than 3,000, and 5 less than 4,000. In one case, the leukocyte level was depressed to 800. A correlation was found between these minimum counts and the doses of individual external irradiation. Similar relationship was found between minimum counts of neutrophils and the dose as shown in Figure 2.




Fig. 2 Correlation between the estimated doses to the whole body and the minimal number of neutrophils.

In most cases a shift to the left of the neutrophils was observed. Staff cells increased in early stage and then gradually decreased. However, in some cases, metamyelocytes and myelocytes appeared in peripheral blood especially with the begin of recovery. In a fatal case who died of liver damage, a considerable number of juvenile neutrophils were observed at the end stage. Lymphopenia was noted between the 2nd and 8th week in all cases. Leukocyte count began to increase after the 8th week. In many cases eosinophilia was remarkable, especially with the indication of recovery. Eosinophilia continued in some cases for several years without proof of parasites, and one case still revealed slight eosinophilia in the 1968 survey.

2. Erythrocytes

At the time of hospitalization a few patients were anemic. No reticulocyte was observed in them at the critical stage. Color index was higher than 1.0. The Price-Jones curves of erythrocyte diameter were displaced to the right of normal at first, and returned to almost normal after one year. These changes corresponded to changes of erythroblast diameters in bone marrow.

3. Platelets, etc.

Platelet counts decreased, reaching a minimum in the 4th to 7th week. Most of the fishermen revealed the values below 100,000/mm³. The results of other tests related to hemorrhage showed depression at the critical stage in general and recovered by the 10th week.

4. Comparison with Other Irradiated Cases

When these blood changes are compared with the changes of other irradiated

to each person differed de-
is cabin.

I was inferred as about 20 to
y in the thyroid region during

found in the urine samples

The radioactivity decreased
in the activity in the urine was

om the fatal case showed that
1, when compared with the
IMURA et al., 1956).

ir content in the urine of the
the same time the results of
rence between fishermen and

ay an important role in early

EARLY STAGE

ns in the early stage.

Fig. 1 Symptoms and signs in
the early stage.

subjects, *i. e.* Hiroshima cases, Marshallese, and cases exposed by reactor accidents (HEMPELMAN et al., 1952; AMANO, 1953; CRONKITE et al., 1956; HASTERLIK and MARINELLI, 1956; GUSKOVA and BAISOCOLOV, 1956; ANDREWS et al., 1961; JAMMET, 1961), it is noted that all these cases similarly revealed the minimal value at about one month after exposure.

5. Bone Marrow

The bone marrow was aplastic at the critical stage in severe cases, which showed remaining and proliferation of plasma cells and reticulum cells. Figure 3 shows the bone marrow of one of the severe cases. Comparing with normal




Fig. 3 Bone marrow of one of severe cases at 5th week.

bone marrow a lack of bone marrow cells was observed. With the beginning of recovery, aplastic marrow changed to hypoplastic and then turned into a type of maturation arrest. In some cases, recovery was not complete even after one year. In less severe cases, the bone marrow was not aplastic but hypoplastic or in a type of maturation arrest even at the critical stage.

6. Morphological and Functional Changes

Several morphological abnormalities were observed: nuclear debris of lymphocytes, binucleated lymphocytes, vacuoles of neutrophils, toxic granulations of neutrophils, degenerative changes of blood cells, mitotic abnormalities of erythroblasts, etc.

Motility and phagocytosis of neutrophils decreased remarkably. After one year, in some cases these functions were still lower than in normal persons.

FOLLOW-UP STUDIES

As above mentioned 16 to 18 fishermen have been examined on an annual basis.

Figure 4 shows the changes of the cumulative distribution curves of leukocytes. The curve of critical stage appeared displaced far to the left of the normal curve, and gradually approached normal values.

However, the curve of the 6th year was still displaced slightly to the left

Fig. 4 Leukocytes of Bik (J) (cumulative).

Fig. 5 Changes in cytes, neutrophils 2

of normal Japa
the curve app
in the cumulat

However,
the changes of
are represent

A small i
of erythrobla
10 years.

Cytogen
results are s

1. The

and cases exposed by reactor ac-
; CRONKITE et al., 1956; HASTERLIK
OV, 1956; ANDREWS et al., 1961;
s similarly revealed the minimal

ical stage in severe cases, which
ells and reticulum cells. Figure
cases. Comparing with normal



at 5th week.

erved. With the begin-
r'astic and then turned
overy was not complete
arrow was not aplastic
n at the critical stage.

ed: nuclear debris of
s, toxic granulations
itotic abnormalities of

markably. After one
in normal persons.

amined on an an-

on curves of leuko-
t the left of the

slightly to the left

Fig. 4 Leukocytes of Bikini patients
(J) (cumulative).





Fig. 5 Changes in the number of leuko-
cytes, neutrophils and lymphocytes.

of normal Japanese. In the case of erythrocytes, 2 years after the exposure, the curve approximately came back to normal. A similar tendency is seen in the cumulative distribution curves of platelets.

However, in a few cases slight neutropenia is still observed. Figure 5 shows the changes of total leukocytes, neutrophils, and lymphocytes of 2 cases which are representative for severe injury.

A small increase of "mitotically connected abnormalities", i. e. karyomeres of erythroblasts, were observed in bone marrow smears of a few cases after 10 years.

Cytogenetical studies have been performed since several years. The results are summarized as follows (ISHIHARA and KUMATORI 1965; 1967; 1969).

1. The frequency of aneuploid cells was 2~3 per cent, which was not

Table 2 Chromosome abnormalities in peripheral lymphocytes of the fishermen, data from 1965 to 1967.

72-h culture:

Cases	No. of cases examined	No. of cells analyzed	Aneuploid cells %	Stable cells % (/)*	unstable cells % (/)*
Control	10	693	2.88	0.14 (1/10)	0.14 (1/10)
Fishermen 1965	13	881	2.05	2.28 (9/13)	0.46 (3/13)
Fishermen 1966	15	1102	2.90	3.45 (14/15)	0.45 (5/15)
Fishermen 1967	16	1248	2.48	2.88 (15/16)	0.56 (6/16)

48-h culture:

Fishermen 1966	10	750	2.27	1.20 (6/10)	0.67 (4/10)
Fishermen 1967	12	950	2.00	2.11 (9/12)	0.74 (6/12)

* Number of cases showing the abnormal cells per total number of cases examined.

so high (Table 2).

2. Unstable abnormalities such as dicentrics, etc. were observed (Table 2).

3. The frequency of stable abnormalities was remarkably higher than in controls (Table 2).

4. The detailed analysis of stable cells revealed that stable abnormalities were divided following 3 types:

- stable cells with balanced chromosome constitution... 67 per cent
- stable cells with deletion of a chromosome... 29 per cent
- stable cells with excessive chromosome material... 4 per cent

5. In 3 cases small chromosomes which are similar to the Ph¹ chromosome were found.

6. The examined fishermen were classified in 3 groups according to the degree of the injuries indicated by the minimum values of neutrophils shown shortly after the exposure. It was found that the mean values of stable and of unstable cells were higher in proportion to the extent of the initial damage. Minimum values of neutrophils roughly corresponded to the estimated external radiation dose.

7. Chromosome abnormalities in bone marrow were observed in 5 cases. All of these abnormalities were limited to stable abnormalities. While stable cells in the peripheral lymphocytes showed various karyotypes of their own, these in the bone marrow were limited to only a few karyotypes.

SPERMATOGENESIS

As a generative tissue, spermatogenesis has many similarities with hematopoiesis. Therefore, the changes of number of spermatozoa is described for reference. The number of spermatozoa decreased, and minimum numbers were obtained in 16 cases about 8 months after the exposure. Lowering of mobility

and morphological abnormalities of recovery were of first signs of regenerative response, with following

The hematological 1954 were reported w

Generally speaking, dose soon after the a sons should be decided findings. From this p to decide the therap

As to the Japan necessary to detect observed in hemato

AMANO, S. (1953):

Gakujutsu Sh

ANDREWS, G. A.,

Diagnosis an

Organization

CRONKITE, E. P.,

Ionizing Rad

Washington

GUSKOVA, A. K. :

Conference

New York.

HASTERLIK, R. J

Conference

New York.

HEMPELMAN, I

Medicine.

ISHIHARA, T. a

ISHIHARA, T.

North Ho

ISHIHARA, T. :

242.

JAMMET, H. F

World I

KIKUCHI, T.,

stitute

KIMURA, K.,

Influen

the Pro

and morphological abnormalities of spermatozoa were also observed. Indications of recovery were observed in some cases after about one year, but mostly first signs of regeneration did not show earlier than about 2 years after exposure, with following production of children.

SUMMARY

The hematological changes of Japanese fishermen exposed to fallout in 1954 were reported with other findings.

Generally speaking, it is difficult to estimate correctly the exposed radiation dose soon after the accident. Therefore, treatment of heavily irradiated persons should be decided by the daily appraisal of clinical and laboratory findings. From this point, hematological changes are an important information to decide the therapy which should be given to heavily irradiated subjects.

As to the Japanese fishermen, it seems to be very important as well as necessary to detect the significance of subtle changes such as they are still observed in hematological and cytogenetical examinations.

REFERENCES

- AMANO, S. (1953): Genshi Bakudan Saigai Chosa Hokokushu. II, 914. Nihon Gakujutsu Shinko Kai, Tokyo.
- ANDREWS, G. A., SITTERSON, B. W., KRETCHMAR, A. L. and BRUCER, M. (1961): Diagnosis and Treatment of Acute Radiation Injury, 27, World Health Organization, Geneva.
- CRONKITE, E. P., BOND, V. P. and DUNHAM, C. H. (ed.) (1956): Some Effects of Ionizing Radiation on Human Beings. U.S. Government Printing Office, Washington, D. C.
- GUSKOVA, A. K. and BAISOGOLOV, G. D. (1956): Proceedings of the International Conference on Peaceful Uses of Atomic Energy, 11, 35, United Nations, New York.
- HASTERLIK, R. J. and MARINELLI, L. D. (1956): Proceedings of the International Conference on Peaceful Uses of Atomic Energy, 11, 25, United Nations, New York.
- HEMPELMAN, L. H., LISCO, H. and HOFFMAN, J. G. (1952): Annals of Internal Medicine, 36, 279.
- ISHIHARA, T. and KUMATORI, T. (1965): Acta Haematologica Japonica, 28, 291.
- ISHIHARA, T. and KUMATORI, T. (1967): Human Radiation Cytogenetics, 144, North Holland Publishing Co., Amsterdam.
- ISHIHARA, T. and KUMATORI, T. (1969): Japan Journal of Genetics, 44, Suppl. 1, 242.
- JAMMET, H. P. (1961): Diagnosis and Treatment of Acute Radiation Injury, 83, World Health Organization, Geneva.
- KIKUCHI, T., WAKISAKA, G., AKAGI, H. and GOTO, H. (1954): Bulletin of the Institute for Chemical Research, 4, Kyoto University, Kyoto.
- KIMURA, K., MINAMI, E., HONDA, M. et al. (1956): Research in the Effects and Influences of the Nuclear Bomb Test Explosions, I, 497, Japan Society for the Promotion of Science, Tokyo.

cytes of the fishermen,

Stable cells % (/)*	unstable cells % (/)*
0.14 (1/10)	0.14 (1/10)
2.28 (9/13)	0.46 (3/13)
3.45 (14/15)	0.45 (5/15)
2.88 (15/16)	0.56 (6/16)
1.20 (8/10)	0.67 (4/10)
2.11 (9/12)	0.74 (6/12)

er of cases examined.

e observed (Table 2).
markably higher than

at stable abnormalities

tion... 67 per cent

20 per cent

... 4 per cent

r to the Ph¹ chromo-

ups according to the
values of neutrophils
mean values of stable
extent of the initial
onded to the estimat-

e observed in 5 cases.
a...ies. While stable
otypes of their own,
aryotypes.

marities with hema-
zoa is described for
mum numbers were
owering of mobility

- KIMURA, K., IKEDA, N., KIMURA, K., KAWANISHI, H. and KIMURA, M. (1956): *ibid.* I, 521.
- KOYAMA, Y., KUMATORI, T. et al. (1955): *Iryo*, 9, 1.
- KUMATORI, T. and MIYOSHI, K. (1963): *Diagnosis and Treatment of Radioactive Poisoning*, 253, International Atomic Energy Agency, Vienna.
- KUMATORI, T., ISHIHARA, T., UEDA, T. and MIYOSHI, K. (1965): *Medical Survey of Japanese Exposed to Fallout Radiation in 1954—A Report after 10 Years*, National Institute of Radiological Sciences, Chiba.
- MIKAMO, Y., MIYOSHI, K., SHIMIZU, K., ISHIKAWA, K., KURIYAMA, S., KOYAMA, Y. and KUMATORI, T. (1956): *Research in the Effects and Influences of the Nuclear Bomb Test Explosions*, II, 1313, Japan Society for the Promotion of Science, Tokyo.
- MIYOSHI, K. and KUMATORI, T. (1955): *Acta Haematologica Japonica*, 18, 379.
- MIYOSHI, K. and KUMATORI, T. (1962): *Proceedings of 8th International Congress of Hematology*, 1, 29, Pan-Pacific Press, Tokyo.
- MIYOSHI, K. and KUMATORI, T. (1964): *Nihon Ketsuekigaku Zensho*, 3, 660, Maruzen Co., Tokyo.

DISCUSSION

Dr. CLIFTON (U.S.A.): Concerning the one fatality, would you care to comment on the nature of the liver damage, and its relationship to external or internal radiation exposure?

Dr. KUMATORI (Japan): Since this fatal case was anemic and revealed bone marrow aplasia, he received blood transfusions. Therefore, serum hepatitis can't be ignored. However, at the same time the existence of radiation-induced liver damage should be considered. At present it is difficult to decide the cause of his liver damage.

We should consider the probability of the radiation-induced liver damage, because in some publications we can see a similar histological picture of the liver of a patient who was irradiated therapeutically.

Dr. SANG (Korea): I thank you Dr. KUMATORI, I would like to ask you something about hemorrhagic tendencies on these cases. Would you tell me how long did hemorrhagic tendencies continue after the irradiation?

Dr. KUMATORI: In my cases the hemorrhagic tendencies were not so severe. The prolongation of the bleeding time was observed in a few cases. You can see normal bleeding time in

most cases. In our cases, we used DUKE's method. The bleeding time is normally 3 minutes, and in 1 or 2 severe cases the bleeding time at the critical stage was as long as about 10 minutes. I think this came back to normal at about 10 weeks or so after the exposures, though I have no correct data here.

Dr. SANG: And how about the fibrinogen concentration?

Dr. KUMATORI: Fibrinogen concentration was almost normal from the beginning of the examination.

Dr. TUBIANA (France): I would like to ask Prof. KUMATORI if he has an idea of how uniform was the dose delivered to your fishermen. I ask you this question because we have a rather large experience of total-body irradiation for mostly kidney transplants in human patients, and there are 2 main differences between your results and the one we have also. The first one is that aplasia of the blood occurs much earlier in our patients, and the rigidity also occurs much earlier. The 2nd one is that the minimum number of leukocytes is much smaller in our cases, being of the order 1 to 200 leukocytes per cubic millimeter, after a dose of the

order of 350 R to 4 and LALANNE, C. M., *logie*, 6, 561, (1963) *Ionizing Radiation (ietic tissue, Int. A (1967))*.

We have compared the curves of Yu cases, and here again same difference, a tion of these data the difference between cases and the patients was that patients the dose with a maximum minus 4 per cent, of accidents the u less, and of the o

Dr. KUMATORI: opinion. In my

M. (1956): *ibid.*

of Radioactive
nna.

Medical Survey of
t after 10 Years,

HOYAMA, Y. and
s of the Nuclear
tion of Science,

onica, 18, 379.
tional Congress

Zensho, 3, 660,

order of 350 R to 400 R (TUBIANA, M. and LALANNE, C. M., *Annales de Radiologie*, 6, 561, (1963) and in *Effects of Ionizing Radiation on the haematopoietic tissue*, Int. At. En. Ag., Vienna (1967)).

We have compared our curves with the curves of Yugoslavian accident cases, and here again we found the same difference, and our interpretation of these data was that probably the difference between the irradiated cases and the purposely irradiated patients was that in the irradiated patients the dose was very uniform, with a maximum variation of plus or minus 4 per cent, whereas in the case of accidents the uniformity was much less, and of the order of 30 per cent.

Dr. KUMATORI: I agree with your opinion. In my cases, I think, the

main irradiation came from externally deposited sources—gamma irradiation. This is a type of uniform irradiation. And this irradiation played an important role in the hematological changes. The fallout material began to precipitate at 7:00 a. m. and continued to fall for about 5 hours. Therefore, the radiation dose to fishermen was at first small and then rapidly increased. As the fishermen began to clean up at about 12:00, the exposure rate decreased thereafter. They were irradiated continuously for 2 weeks with decreasing dose rates. Aside from this external irradiation, internal exposure may have had some effects. But the internal irradiation seemed to have only a slight effect on the acute hematological changes.

cases, we used
bleeding time is
and in 1 or 2
ng time at the
ng as about 10
came back to
or so after
I have no

about the fib-

ogen concent-
mal from the
tion.

I would like
if he has an
was the dose
ermen. I ask
se we have a
total-body
kidney trans-
s, and there
etween your
e have also.
lasia of the
rlier in our
also occurs
ne is that the
eukocytes is
ses, being of
kocytes per
dose of the